

REDISCOVERY AND INTEGRATIVE IDENTIFICATION OF *MILAX GAGATES* FROM JAPAN (GASTROPODA: MILACIDAE)

TAICHI KATSUMI¹, NAOTO SAWADA^{2*}

¹School of Science, Osaka Metropolitan University, Japan

²School of Science, The University of Tokyo, Japan (e-mail: sawada@zoo.zool.kyoto-u.ac.jp)

* corresponding author

ABSTRACT: *Milax gagates*, native to the western Mediterranean region, has been introduced to many countries including Japan. Despite the need for an anatomical examination for its accurate identification, records of this species in Japan have been based on brief lists in the 20th century and its recent distribution remains unclear. In this study, we rediscovered slugs of *M. gagates* from Tokyo, Japan, and identified them by examining anatomical traits and a mitochondrial cytochrome c oxidase subunit I (COI) haplotype. This species is believed to be established in the study area based on field observations in 2022 and 2024. The haplotype of the Japanese specimens corresponded to those of non-native populations in the UK, the Netherlands, Portugal, and the Iberian Peninsula, suggesting that it might not only have been originated from the native range, but also from secondary introduction from non-native areas.

KEY WORDS: DNA barcoding; exotic species; genitalia; haplotype network

INTRODUCTION

Anthropogenic introductions of invasive species have significantly impacted native ecosystems, economic activity, and human health (LOWE et al. 2000, SIMBERLOFF et al. 2005, DIDHAM et al. 2007, PYŠEK et al. 2020). Invasive slugs reduce the populations of native terrestrial molluscs by decreasing their reproductive rates and increasing mortality of native species, as well as increasing hybridisation among them (ROLLO 1983, HATTELAND et al. 2015, REISE et al. 2020). Slugs also cause economic and sanitary damage to their non-native areas by plant feeding (HERBERT 2010, ROWSON et al. 2014b, LE GALL & TOOKER 2017) and parasite transmission (HOLLINGSWORTH et al. 2013). Thus, investigating their current distribution, activity patterns, and control methods are important to reduce their long-term risks (GALVIS & MORENO 2017, MORII et al. 2018, NURINSIYAH & HAUSDORF 2019, VENDETTI et al. 2019).

Milax gagates (Draparnaud, 1801) is a slug species with a wide non-native distribution. This species is

assumed to be native to the western Mediterranean region, while it has been recorded in many countries in Europe, North and South America, and Oceania (LOVETT & BLACK 1920, WIKTOR 1987, HERBERT 2010, THOMAS et al. 2010). This species has been confused with congeners, such as *M. nigricans* (Philippi, 1836), because of their similar appearance, but their genital characteristics proved to be useful in distinguishing the species from other Milacidae species (WIKTOR 1987, HUTCHINSON & REISE 2013, ROWSON et al. 2014b). Recently, these two species have been identified by DNA barcoding using mitochondrial cytochrome c oxidase subunit I (COI) fragments (TURÓCI et al. 2023).

The non-native distribution of *M. gagates* includes Japan (WIKTOR 1987, BARKER 1999, HERBERT 2010). This species was reported from Tokyo (YAMAGUCHI & HABE 1955) and Kanagawa Prefecture (KEIO UHNSG 1991) in eastern Japan, and has been listed as an alien species in Japan (KURODA 1963, AZUMA 1982, MINATO 1988, SUBCOMMITTEE ISISGWPM

2002, CHIBA PREFECTURAL GOVERNMENT 2020). Field records of *M. gagates* are limited to these two prefectures in Japan, and the species has only been found during phytosanitary inspections at airports in recent years (MATSUMOTO & KUROSUMI 2004). However, proper identification and morphological descriptions were not provided by the two brief field records, and to our knowledge, neither anatomical nor genetic examinations have been attempted on

Japanese *Milax* specimens. Therefore, the recent establishment status and identity of *Milax* species have not yet been clarified in Japan.

We recently rediscovered *Milax* slugs on a river bank in Tokyo, Japan, and conducted species identification and habitat surveys, aiming to reveal the true identity of these milacid slugs and to find out whether they are established in Japan or remained single introductions.

MATERIAL AND METHODS

FIELD SURVEY AND SPECIMEN PREPARATION

A field survey of slugs was conducted on the river-side of the Edogawa River in Shibamata, Katsushika-ku, Tokyo Metropolitan, Japan ($35^{\circ}45'25''\text{N}$, $139^{\circ}52'54''\text{E}$) on 14 May 2022, 17 April 2024, and 10 May 2024 (Fig. 1). The dominant vegetation of the habitat, rough abundance and habitat range of slugs, and sympatric terrestrial molluscs were visually investigated on 10 May 2024. The molluscs were identified following AZUMA (1982), MINATO (1989), HWANG et al. (2021), and MOLLUSCABASE (2024).

Among *Milax* slugs observed on the three survey days, two specimens collected on 14 May 2022 and 14 specimens collected on 10 May 2024, were used for morphological and/or genetic analyses in this study. They were stored at 13°C in a refrigerator until further analysis. The specimens were prepared and photographed according to the method described by TURÓCI et al. (2023). After the morphological and genetic analyses, the specimens were deposited at the Zoological Collection of Kyoto University (KUZ Z5701–Z5707).




Fig. 1. Map representing collection locality of *Milax gagates* in Tokyo, Japan. The map was drawn based on data from JARVIS et al. (2008), MINISTRY LITTIJ (2020), and GEOSPATIAL INFORMATION (2024)

MORPHOLOGICAL EXAMINATION

The 14 slugs collected on 10 May 2024 were examined. The external morphology of the living and preserved specimens was photographed using a D7500 camera (Nikon Corporation, Tokyo, Japan) with a Nikon-compatible SP 90 mm f/2.8 1:1 macro lens (Tamron Co., Ltd., Saitama, Japan). The specimens were dissected using a YS05T stereomicroscope (Micronet Inc., Kawaguchi, Japan). The reproductive organs were photographed using a Swiftcam SC1003-CK camera (SwiftCam Technologies Group Co., Ltd., Kwai Chung, Hong Kong) attached to the stereomicroscope. Based on the morphological distinctiveness of *Milax* species indicated previously (WIKTOR 1987, HUTCHINSON & REISE 2013, TURÓCI et al. 2023), we focused on the shapes of the atrial stimulator. In addition, position of the vas deferens outlet, shape and size of the bursa copulatrix, and body colour, which has been identified as less reliable (HUTCHINSON & REISE 2013, TURÓCI et al. 2023), was also investigated. After dissection, the shells and radulae were extracted from six and five specimens, respectively, using 1 M sodium hydroxide. According to GERMAIN (1930), WIKTOR (1987), and ROWSON et al. (2014b), shell and radula morphology between the previous observations and the present specimens were compared. The body length of the preserved specimen and shell length were measured using ImageJ version 1.51 (SCHNEIDER et al. 2012). The extracted radulae were

photographed using a JSM-6510LV scanning electron microscope (JEOL Ltd., Tokyo, Japan) after platinum coating using a JEC-3000FC (JEOL Ltd.).

GENETIC ANALYSIS

Morphological identification of the dissected specimens was confirmed using a partial region of mitochondrial COI. Genomic DNA was extracted from two specimens collected on 14 May 2022 and four specimens collected on 10 May 2024, following the procedure outlined by OKAMOTO et al. (2006). Polymerase chain reaction and nucleotide sequencing with the methods by SAWADA et al. (2021) using the primers LCO1490 (5'-GGTCAACAAATCATAAAGATATTGG-3') and HCO2198 (5'-TAAACTTCAGGGTGACCAAAAAATCA-3') (FOLMER et al. 1994). The newly obtained sequences were deposited in the International Nucleotide Sequence Database (INSD) through the DNA Databank of Japan (LC850668–LC850673). Genetic variations of the COI sequence were visually evaluated using MEGA version 11 (TAMURA et al. 2021) against 20 sequences of *M. gagates* (including sequences labelled as *Milax* sp.) and *M. nigricans* provided by ROWSON et al. (2014a), GÓMEZ-RODRÍGUEZ et al. (2019), and TURÓCI et al. (2023). In addition, their haplotype relationships were inferred by statistical parsimony (CLEMENT et al. 2000) using POPART v1.7 (LEIGH & BRYANT 2015).

RESULTS

IDENTIFICATION

Slugs collected from the banks of the Edogawa River had light to dark brownish grey bodies with

a distinct middorsal keel on the posterior mantle edge to the tip of the tale (Figs 2–7). The sole was mostly brownish light grey, and the dark grey pneumostome rim was conspicuous on the lighter-bodied



Figs 2–7. Dorsal, lateral, and ventral views of living specimens of *Milax gagates* collected from Tokyo, Japan: 2–4 – KUZ Z5701; 5–7 – KUZ Z5702. Slugs were stored at 13 °C in a refrigerator for three days from collection to being photographed


Table 1. List of examined Japanese *Milax gagates* specimens with INSD accessions, body measurements, and anatomical observations. Two specimens, MGJ15 and MGJ16, were collected in 2022 and the others were collected in 2024.

Specimen # (voucher #)	INSD accession	Life- stage	Body length (preserved) (mm)	Body colour	Rim of pneu- mostome	Shell length (mm)	Shape of atrial stimulator	Papillae of stimulator	Shape of bursa copu- latrix	Shape of bursa trunk	Connection of vas deferens to epiphallus
MGJ01 (KUZ Z5701)	LC850670	adult	25.9	dark grey	indistinct	3.5	short, tapered, straight	small papillae from middle to periphery of tip of stimulator	large, oval	short, thick	slightly displaced at apex
MGJ02 (KUZ Z5702)	LC850671	adult	22.2	brownish light grey	distinct	2.9	short, tapered, slightly curved	small papillae on periphery of tip of stimulator	large, oval	short, thick	slightly displaced at apex
MGJ03 (KUZ Z5703)	LC850672	adult	23.9	brownish light grey	distinct	3.7	long, tapered, strongly curved	small papillae on periphery of tip of stimulator	large, oval	short, thick	slightly displaced at apex
MGJ04 (KUZ Z5704)	LC850673	adult	26.4	dark grey	indistinct	4.1	short, tapered, strongly curved	indistinct	large, oval	long, narrow	slightly displaced at apex
MGJ05 (KUZ Z5705)		adult	20.7	brownish light grey	distinct	3.2	long, tapered, strongly curved	small papillae from middle to periphery of tip of stimulator	small, oval	long, narrow	slightly displaced at apex
MGJ06 (KUZ Z5705)		adult	27.5	dark grey	indistinct		short, tapered, straight	small papillae from middle to periphery of tip of stimulator	large, oval	short, thick	slightly displaced at apex
MGJ07 (KUZ Z5705)		adult	25.9	dark grey	indistinct	4.3	long, tapered, slightly curved	small papillae from middle to periphery of tip of stimulator	small, oval	short, thick	slightly displaced at apex
MGJ08 (KUZ Z5705)		adult	24.9	dark grey	indistinct		long, tapered, strongly curved	small papillae from middle to periphery of tip of stimulator	small, oval	long, narrow	slightly displaced at apex
MGJ09 (KUZ Z5705)		adult	19.4	dark grey	indistinct		short, tapered, straight	small papillae from middle to periphery of tip of stimulator	large, oval	short, thick	slightly displaced at apex
MGJ10 (KUZ Z5705)		adult	17.3	dark grey	indistinct		long, tapered, slightly curved	small papillae from middle to periphery of tip of stimulator	large, elon- gated	short, thick	slightly displaced at apex
MGJ11 (KUZ Z5705)		adult	18.3	dark grey	indistinct		long, tapered, straight	small papillae on periphery of tip of stimulator	large, elon- gated	short, thick	slightly displaced at apex
MGJ12 (KUZ Z5705)		adult	19.1	dark grey	indistinct		long, tapered, straight	small papillae from middle to periphery of tip of stimulator	large, oval	short, thick	at apex
MGJ13 (KUZ Z5705)		adult	20.2	dark grey	indistinct		long, tapered, strongly curved	small papillae on periphery of tip of stimulator	large, oval	short, thick	slightly displaced at apex
MGJ14 (KUZ Z5705)		juve- nile	19.8	dark grey	indistinct		immature	immature	small, oval	long, narrow	slightly displaced at apex
MGJ15 (KUZ Z5706)	LC850668	juve- nile		brownish light grey							
MGJ16 (KUZ Z5707)	LC850669	juve- nile		brownish light grey							

specimens. Well-developed genitalia were observed in 13 specimens, whereas they were immature in one specimen (Table 1). In mature specimens, the atrial stimulator in the genitalia was short to long, tapered, and almost straight or strongly curved at the tip (Figs 8–11). Three to four small papillae were observed from the middle to the periphery of the tip of the stimulator except in one specimen which possessed a stimulator without distinct papillae

(Table 1). Most specimens had an oval, large bursa copulatrix, a short, thick bursa trunk, and vas deferens connected slightly displaced at the apex of the epiphallus. They possessed a 2.9–4.3 mm oval, thin shell inside the anterior dorsal mantle (Fig. 12).

The radula possessed one rachidian and 32–44 lateral and marginal teeth on each side of the transverse row (Figs 13 & 14). The rachidian and lateral teeth had one arrow-headed large mesocone and one



Figs 8–14. Reproductive organs (8–11), shell (12), and radula (13–14) of *Milax gagates* collected from Tokyo, Japan: 8 – entire genitalia, KUZ Z5701; 9 – genitalia around atrium, KUZ Z5702; 10–11 – atrial stimulator, KUZ Z5701 (10), Z5702 (11); 12 – KUZ Z5705; 13 – rachidian and lateral teeth, KUZ Z5702; 14 – lateral and marginal teeth, KUZ Z5702. Slugs were stored at 13 °C in a refrigerator for three to four days from collection to being dissected. Arrows indicate papillae on the atrial stimulator. Abbreviations: atg – atrial gland; ag – albumen gland; bc – bursa copulatrix; epi – epiphallus; hd – hermaphroditic duct; hg – hermaphroditic gland; ovi – oviduct; p – penis; so – spermoviduct; sti – atrial stimulator; vd – vas deferens. Scale bars 5 mm (8), 1 mm (9, 12), 0.5 mm (10, 11), 0.1 mm (13, 14)

smaller ectocone and endocone on both sides. The mesocone was approximately 3.5 times longer than the ectocone and endocone in the inner rows of the radula, and these minor cusps were smaller in the outer rows. The marginal tooth possessed one pointed mesocone and was smaller in the outer rows.

The 655 bp COI sequences obtained from six specimens showed the identical haplotype. These sequences matched completely in aligned positions with those of *M. gagates* from the UK (KF894243) and the Netherlands and Portugal (OP663302), and *Milax* sp. from the Iberian Peninsula (MF983226). The statistical parsimony network calculated based on the 12 haplotypes separated the sequences into two major groups by 21 mutational steps (Fig. 15). The network also showed that sequences obtained from the specimens of *M. gagates* (orange) and *M. nigricans* (red), which were identified with morphological traits, were divided at least in 26 mutational steps.

HABITAT

Slugs were observed on paved roads and grassy areas at the same location on the banks of the Edogawa River on the three survey days. The distribution of the slugs was restricted to an area of approximately 0.3 ha and more than 150 *Milax* slugs with various body sizes were observed. The highest densities of slugs were found in sunny areas with short bushes, where herbaceous plants such as *Avena fatua*, *Trifolium dubium*, and *Vicia sativa nigra* were dominant. *Milax* slugs were observed sympatrically with the following five terrestrial mollusc species: *Granulilimax fuscicornis* Minato, 1989, *Allopeas clavulinum kyotoense* (Pilsbry et Hirase, 1904), *Ambigolimax valentianus* (Férussac, 1821), *Acusta* cf. *sieboldiana* (Pfeiffer, 1850), *Bradybaena similaris* (Férussac, 1822).




Fig. 15. Statistical parsimony network showing the relationships between the 12 haplotypes detected in a partial COI sequence of *Milax* species. Each connection indicates a single mutation, and hatch marks represent missing intermediate haplotypes. Sequences obtained from specimens identified with anatomical traits were coloured: *M. gagates* (orange) and *M. nigricans* (red)

DISCUSSION

In the present study, slugs collected from Japan were identified as *M. gagates* both morphologically and genetically. The anatomical characteristics observed in most Japanese specimens were consistent with those of *M. gagates* identified in previous studies (WIKTOR 1987, HUTCHINSON & REISE 2013). Particularly, the papillae of the atrial stimulator were

small or indistinct in all specimens, which is the reliable character for discriminating between *M. gagates* and *M. nigricans* (HUTCHINSON & REISE 2013). On the other hand, significant variations were observed in less diagnostic vas deferens, bursa copulatrix, bursa trunk, and body colour. The shell and radula morphology of Japanese slugs was similar to those



of Milacidae species (GERMAIN 1930, WIKTOR 1987, ROWSON et al. 2014b). The identical COI haplotype of the six Japanese slugs were the same as or close to those of *M. gagates* identified in previous studies (ROWSON et al. 2014a, TURÓCI et al. 2023).

The haplotype network separated the 26 sequences into two major groups. The sequences of *M. gagates* and *M. nigricans* identified in the previous studies using anatomical traits belonged to different groups. The group of *M. gagates* included the sequences of *M. gagates* from UK, Ireland, Netherlands, Portugal, and Belgium (ROWSON et al. 2014a, TURÓCI et al. 2023), *Milax* sp. from Iberian Peninsula (GÓMEZ-RODRÍGUEZ et al. 2019), and the Japanese *Milax* obtained in this study. Furthermore, the haplotype of the Japanese specimens was identical to those from the UK, the Netherlands, Portugal, and the Iberian Peninsula. The native range of *M. gagates* has been estimated to be the coastal and insular regions of the western Mediterranean, and the species has artificially expanded into the regions where the same haplotypes as those in Japan have been obtained (LOVETT & BLACK 1920, WIKTOR 1987). Because all three sequences identical to Japanese ones were obtained outside the native range of *M. gagates* (WIKTOR 1987), it is possible that the Japanese population was formed by secondary introduction from its non-native distribution, as well as direct transportation from its origin.

In the present study, *M. gagates* were found in the same locality in 2022 and 2024, and 150 mature and immature slugs were observed in 2024. Given that a twice-a-year reproduction has been revealed in this species (WIKTOR 1987), it has likely established in this humid subtropical area in Japan.

Japanese *Milax* was observed with five terrestrial mollusc species. Among them, *G. fuscicornis*, *Al. clavulinum kyotoense*, and *Ac. cf. sieboldiana*, are assumed

native to Japan and *Am. valentianus* are thought to be non-native (AZUMA 1982, MINATO 1989, HWANG et al. 2021). There is still potential for both native and non-native status for *B. similaris* in Japan (HIRANO et al. 2023). The introduction of alien slugs has contributed to the decline of native land snails through parasitic infections and predation (HATTELAND et al. 2013, HOWE et al. 2020). Feeding damage to crops and rare indigenous plants, and the spread of their diseases have also been partially attributed to non-native slugs, including *M. gagates* (HERBERT 2010). Although the ecological impact of the Japanese *M. gagates* could not be evaluated in this study, they can contribute to the decline in the mollusc and plant species at the survey sites. Therefore, further investigation is required to evaluate its impact on native ecosystems and its expansion.

ACKNOWLEDGEMENTS

The authors are grateful to TAKUTO MIYAI for his assistance with the field surveys and sample collection. We also thank TATSUKI NISHIOKA for his support in identifying plant species. We also express our gratitude to TAIJI KUROZUMI, HIROSHI FUKUDA (Okayama University), TOMOTSUGU ARIKAWA (Keio University), and KINJI FUKUYAMA for providing information on Japanese *M. gagates*. The authors are grateful to REI UESHIMA (The University of Tokyo) for assisting with radular observation. We also express our appreciation to TAKAFUMI NAKANO (Kyoto University) for his assistance with specimen deposition at the KUZ. We are grateful to three anonymous reviewers for their constructive comments. This study was partly supported by JSPS KAKENHI (Grant No. JP24KJ0045). We thank Editage (www.editage.jp) for English language editing.

REFERENCES

- AZUMA M. 1982. Colored illustrations of the land snails of Japan. Hoikusha Publishing, Osaka. [in Japanese]
- BARKER G. M. 1999. Naturalised terrestrial Stylommatophora (Mollusca: Gastropoda). Fauna of New Zealand 38: 1–253.
- CHIBA PREFECTURAL GOVERNMENT 2020. List of invasive species in Chiba Prefecture. Nature Conservation Division, Environmental and Community Affairs Department, Chiba Prefectural Government, Chiba. [in Japanese]
- CLEMENT M., POSADA D., CRANDALL K. A. 2000. TCS: a computer program to estimate gene genealogies. Molecular Ecology 9: 1657–1659. <https://doi.org/10.1046/j.1365-294x.2000.01020.x>
- DIDHAM R. K., TYLIANAKIS J. M., GEMMELL N. J., RAND T. A., EWERS R. M. 2007. Interactive effects of habitat modification and species invasion on native species decline. Trends in Ecology and Evolution 22: 489–496. <https://doi.org/10.1016/j.tree.2007.07.001>
- FOLMER O., BLACK M., HOEH W., LUTZ R., VRIJENHOEK R. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. Molecular Marine Biology and Biotechnology 3: 294–299.
- GALVIS F., MORENO L. 2017. Identification of Cry1 genes in *Bacillus thuringiensis* isolates and their toxic effect against *Milax gagates*, a pest on lettuce (*Lactuca sativa*). Revista Chapingo Serie Horticultura 24: 97–106. <https://doi.org/10.5154/rchsh.2017.07.025>

- GEOSPATIAL INFORMATION 2024. Goespatial Information Authority of Japan: GSI maps. Available online at <https://maps.gsi.go.jp/> [accessed on 10 November 2024]
- GERMAIN L. 1930. Mollusques terrestres et fluviatiles (première partie). *Faune de France* 21: 1–477.
- GÓMEZ-RODRÍGUEZ C., MILLER K. E., CASTILLEJO J., IGLESIAS-PIÑEIRO J., BASELGA A. 2019. Understanding dispersal limitation through the assessment of diversity patterns across phylogenetic scales below the species level. *Global Ecology and Biogeography* 28: 353–364. <https://doi.org/10.1111/geb.12857>
- HATTELAND B. A., ROTH S., ANDERSEN A., KAASA K., STØA B., SOLHØY T. 2013. Distribution and spread of the invasive slug *Arion vulgaris* Moquin-Tandon in Norway. *Fauna Norvegica* 32: 13–26. <https://doi.org/10.5324/fn.v32i0.1473>
- HATTELAND B. A., SOLHØY T., SCHANDER C., SKAGE M., PROSCHWITZ T. VON, NOBLE L. R. 2015. Introgression and differentiation of the invasive slug *Arion vulgaris* from native *A. ater*. *Malacologia* 58: 303–321. <https://doi.org/10.4002/040.058.0210>
- HERBERT D. G. 2010. The introduced terrestrial Mollusca of South Africa. SANBI Biodiversity Series 15. South African National Biodiversity Institute, Pretoria.
- HIRANO T., YAMAZAKI D., ITO S., SATO M. P., MATSUO A., SAITO T., NISHI H., YE B., DONG Z., TU D. V., SHAU-HWAI A. T., SUYAMA Y., CHIBA S. 2023. Reconsidering invasion history of common land snails in Japan through genome-wide analyses. *Biological Invasions* 25: 3535–3549. <https://doi.org/10.1007/s10530-023-03123-2>
- HOLLINGSWORTH R. G., HOWE K. B., JARVI S. I. 2013. Control measures for slug and snail hosts of *Angiostrongylus cantonensis*, with special reference to the semi-slug *Parmarion martensi*. *Hawai'i Journal of Medicine & Public Health* 72: 75–80.
- HOWE D. K., HA A. D., COLTON A., LEY I. T. DE, RAE R. G., ROSS J., WILSON M., NERMUT J., ZHAO Z., MC DONNELL R. J., DENVER D. R. 2020. Phylogenetic evidence for the invasion of a commercialized European *Phasmarhabditis hermaphrodita* lineage into North America and New Zealand. *PLoS ONE* 15: e0237249. <https://doi.org/10.1371/journal.pone.0237249>
- HUTCHINSON J. M. C., REISE H. 2013. A persisting population of an introduced slug, *Milax nigricans*, in Dunkirk, France. *Mitteilungen der Deutschen Malakozoologischen Gesellschaft* 89: 35–38.
- HWANG C. C., ZHOU W. C., GER M. J., GUO Y., QIAN Z. X., WANG Y. C., TSAI C. L., WU S. P. 2021. Biogeography of land snail genus *Acusta* (Gastropoda: Camaenidae): Diversification on East Asian islands. *Molecular Phylogenetics and Evolution* 155: 106999. <https://doi.org/10.1016/j.ympev.2020.106999>
- JARVIS A., REUTER H. I., NELSON A., GUEVARA E. 2008. Hole-filled seamless SRTM data V4. International Centre for Tropical Agriculture (CIAT). Available online at <https://srtm.csi.cgiar.org/> (accessed 10 November 2024).
- KEIO UHNSG 1991. Report on the nature survey of the Keio University Hiyoshi Campus area. Keio University Hiyoshi Nature Survey Group, Yokohama. [in Japanese]
- KURODA T. 1963. A catalogue of the non-marine mollusks of Japan, including the Okinawa and Ogasawara Islands. The Malacological Society of Japan, Tokyo.
- LE GALL M., TOOKER J. F. 2017. Developing ecologically based pest management programs for terrestrial molluscs in field and forage crops. *Journal of Pest Science* 90: 825–838. <https://doi.org/10.1007/s10340-017-0858-8>
- LEIGH J. W., BRYANT D. 2015. POPART: full-feature software for haplotype network construction. *Methods in Ecology and Evolution* 6: 1110–1116. <https://doi.org/10.1111/2041-210X.12410>
- LOVETT A. L., BLACK A. B. 1920. Station bulletin 170. The gray garden slug: with notes on allied forms. Oregon Agricultural College Experiment Station, Oregon.
- LOWE S., BROWNE M., BOUDJELAS S., DE POORTER M. 2000. 100 of the world's worst invasive alien species: a selection from the global invasive species database. Invasive Species Specialist Group, Auckland.
- MATSUMOTO N., KUROZUMI T. 2004. A list of slugs (Gastropoda: Pulmonata) intercepted at plant quarantine in Japan. *Research Bulletin of the Plant Protection Service* 40: 127–133.
- MINATO H. 1988. A systematic and bibliographic list of the Japanese land snails. Publication Committee of a Systematic and Bibliographic List of the Japanese Land Snails, Shirahama [in Japanese]
- MINATO H. 1989. A remarkable slug, *Granulilimax fuscicornis* n. gen. et sp. from Japan. *Venus* 48: 255–258. https://doi.org/10.18941/venusjtm.48.4_255
- MINISTRY LITJ 2020. Ministry of Land, Infrastructure, Transport and Tourism of Japan: Lake data. 2005 version. Available online at <https://nlftp.mlit.go.jp/ksj/gml/datalist/KsjTmplt-W09-2005.html> [in Japanese] (accessed 10 November 2024)
- MOLLUSCABASE 2024. MolluscaBase (eds). Available online at <https://www.molluscabase.org> (accessed 10 November 2024)
- MORII Y., OHKUBO Y., WATANABE S. 2018. Activity of invasive slug *Limax maximus* in relation to climate conditions based on citizen's observations and novel regularization based statistical approaches. *Science of the Total Environment* 637–638: 1061–1068. <https://doi.org/10.1016/j.scitotenv.2018.04.403>
- NURINSIYAH A. S., HAUSDORF B. 2019. Listing, impact assessment and prioritization of introduced land snail and slug species in Indonesia. *Journal of Molluscan Studies* 85: 172–176. <https://doi.org/10.1093/mollus/eyy062>
- OKAMOTO T., MOTOKAWA J., TODA M., HIKIDA T. 2006. Parapatric distribution of the lizards *Plestiodon* (formerly *Eumeces*) *laticutatus* and *P. japonicus* (Reptilia: Scincidae) around the Izu Peninsula, central Japan, and its biogeographic implications. *Zoological Science* 23: 419–425. <https://doi.org/10.2108/zsj.23.419>



- PYŠEK P., HULME P. E., SIMBERLOFF D., BACHER S., BLACKBURN T. M., CARLTON J. T., DAWSON W., ESSL F., FOXCROFT L. C., GENOVESI P., JESCHKE J. M., KÜHN I., LIEBHOLD A. M., MANDRAK N. E., MEYERSON L. A., PAUCHARD A., PERGL J., ROY H. E., SEEBENS H., VAN KLEUNEN M., VILÀ M., WINGFIELD M. J., RICHARDSON D. M. 2020. Scientists' warning on invasive alien species. *Biological Reviews* 95: 1511–1534.
<https://doi.org/10.1111/brv.12627>
- REISE H., SCHWARZER A., HUTCHINSON J. M. C., SCHLITT B. 2020. Genital morphology differentiates three subspecies of the terrestrial slug *Arion ater* (Linnæus, 1758) s.l. and reveals a continuum of intermediates with the invasive *A. vulgaris* Moquin-Tandon, 1855. *Folia Malacologica* 28: 1–34.
<https://doi.org/10.12657/folmal.028.001>
- ROLLO C. D. 1983. Consequences of competition on the reproduction and mortality of three species of terrestrial slugs. *Population Ecology* 25: 20–43.
<https://doi.org/10.1007/BF02528782>
- ROWSON B., ANDERSON R., TURNER J. A., SYMONDSON W. O. C. 2014a. The slugs of Britain and Ireland: undetected and undescribed species increase a well-studied, economically important fauna by more than 20%. *PLOS ONE* 9: e91907.
<https://doi.org/10.1371/journal.pone.0091907>
- ROWSON B., TURNER J., ANDERSON R., SYMONDSON B. 2014b. Slugs of Britain and Ireland. Identification, understanding and control. FSC / National Museum of Wales, Telford.
- SAWADA N., HWANG C. C., HARL J., NAKANO T. 2021. Integrative taxonomy of insular land snails of the genus *Sicradiscus* Páll-Gergely, 2013 (Gastropoda, Plectopylidae) with description of a new species. *Journal of Zoological Systematics and Evolutionary Research* 59: 1798–1815.
<https://doi.org/10.1111/jzs.12553>
- SCHNEIDER C. A., RASBAND W. S., ELICEIRI K. W. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* 9: 617–675.
<https://doi.org/10.1038/nmeth.2089>
- SIMBERLOFF D., PARKER I. M., WINDLE P. N. 2005. Introduced species policy, management, and future research needs. *Frontiers in Ecology and the Environment* 3: 12–20.
[https://doi.org/10.1890/1540-9295\(2005\)003\[0012:ispmaf\]2.0.co;2](https://doi.org/10.1890/1540-9295(2005)003[0012:ispmaf]2.0.co;2)
- SUBCOMMITTEE ISISGWPM 2002. Subcommittee on Invasive Species Issues of the Study Group on Wildlife Protection Measures: Policy for dealing with invasive species (non-native species). Available online at <https://www.env.go.jp/content/900493508.pdf> (accessed 10 November 2024). [in Japanese]
- TAMURA K., STECHER G., KUMAR S. 2021. MEGA11: Molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution* 38: 3022–3027.
<https://doi.org/10.1093/molbev/msab120>
- THOMAS A., MC DONNELL R., HARWOOD J. 2010. A field guide to the slugs of Kentucky. University of Kentucky, Lexington.
- TURÓCI Á., HUTCHINSON J. M. C., SCHLITT B., REISE H., RAPALA M., PÁLL-GERGELY B. 2023. Five new introduced terrestrial slugs in Hungary. *BioInvasions Records* 12: 711–729.
<https://doi.org/10.3391/bir.2023.12.3.08>
- VENDETTI J. E., BURNETT E., CARLTON L., CURRAN A. T., LEE C., MATSUMOTO R., MC DONNELL R., REICH I., WILLADSEN, O. 2019. The introduced terrestrial slugs *Ambigolimax nyctelius* (Bourguignat, 1861) and *Ambigolimax valentianus* (Férussac, 1821) (Gastropoda: Limacidae) in California, with a discussion of taxonomy, systematics, and discovery by citizen science. *Journal of Natural History* 53: 1607–1632.
<https://doi.org/10.1080/00222933.2018.1536230>
- WIKTOR A. 1987. Milacidae (Gastropoda, Pulmonata) - systematic monograph. *Annales Zoologici* 41: 202–206.
<https://rcin.org.pl/dlibra/publication/72987/edition/>
- YAMAGUCHI N., HABE T. 1955. Studies on the Japanese slugs (1). *Venus* 18: 234–240.
https://doi.org/10.18941/venusjjmc.18.4_234

Received: November 15th, 2024

Revised: January 12th, 2025

Accepted: January 18th, 2025

Published on-line: March 4th, 2025

